

AD-A127 031

ELECTRONICALLY INDUCED PHASE TRANSFORMATIONS(U)
NORTHWESTERN UNIV EVANSTON IL DEPT OF PHYSICS AND
ASTRONOMY A J FREEMAN 30 NOV 82 AFOSR-TR-83-0263

1/1


UNCLASSIFIED

AFOSR-81-0024

fig 7/4 N6



END
DATE
FILMED
5 - H3
DTIC



MICROCOPY RESOLUTION TEST CHART
NATIONAL BUREAU OF STANDARDS-1963-A

Unclassified

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

(6)

REPORT DOCUMENTATION PAGE		READ INSTRUCTIONS BEFORE COMPLETING FORM	
1. REPORT NUMBER AFOSR-TR- 83 - 0263	2. GOVT ACCESSION NO. ADA127031	3. RECIPIENT'S CATALOG NUMBER	
4. TITLE (and Subtitle) ELECTRONICALLY INDUCED PHASE TRANSFORMATIONS		5. TYPE OF REPORT & PERIOD COVERED Abstract Oct. 1, 1981-Sept. 30, 1982	
6. AUTHOR(s) Arthur J. Freeman		7. PERFORMING ORG. REPORT NUMBER	
8. CONTRACT OR GRANT NUMBER(s) AFOSR-81-0024		9. PROGRAM ELEMENT, PROJECT, TASK AREA & WORK UNIT NUMBERS 61102F 2301-1A8	
10. CONTROLLING OFFICE NAME AND ADDRESS AFOSR/NP Bolling AFB, Bldg. 410 Washington, DC 20332		11. REPORT DATE 30 November 1982	
12. MONITORING AGENCY NAME & ADDRESS (if different from Controlling Office)		13. NUMBER OF PAGES 9	
14. DISTRIBUTION STATEMENT (of this Report) Approved for public release; distribution unlimited.		15. SECURITY CLASS. (of this report) Unclassified	
16. DISTRIBUTION STATEMENT (of the abstract entered in Block 20, if different from Report)		17. SUPPLEMENTARY NOTES	
18. KEY WORDS (Continue on reverse side if necessary and identify by block number) electronically induced phase transformations high field superconductivity in Chevrel phase materials soft phonon modes and superconductivity in C15 compounds		19. ABSTRACT (Continue on reverse side if necessary and identify by block number) Our recent development of self-consistent local (spin) density energy band approaches have provided a powerful theoretical/computational tool for determining the electronic structure and properties of complex materials. The importance of charge transfer between constituent atom species has been demonstrated as has its inclusion by means of accurate self-consistent solutions. Some examples of areas in which progress has been achieved include: the high field superconductivity in Chevrel phase compounds; the unusual →	

AD A 127031

DTIC FILE COPY

DTIC
ELECTE
S APR 21 1983 D
E

DD FORM 1473 JAN 73

EDITION OF 1 NOV 65 IS OBSOLETE

Unclassified

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

magnetic and superconducting properties of some C15 compounds including a better understanding of their relationship between the electronic, lattice and superconducting properties of these materials; self-consistent energy band calculations including all electrons and all atoms in the 16 atoms per unit cell of the linear chain transition metal trichalcogenide, TaSe_3 ; and a detailed assessment of theoretical determinations of the electron phonon coupling parameter in metals and intermetallic compounds.

Accession For	
NTIS GRA&I	<input checked="checked" type="checkbox"/>
DTIC TAB	<input type="checkbox"/>
Unannounced	<input type="checkbox"/>
Justification	
By	
Distribution/	
Availability Codes	
Dist	Avail and/or Special
A	



PROGRESS REPORT

Our recent development of self-consistent local (spin) density energy band approaches to determining the electronic structure and properties of complex materials have provided a powerful theoretical/computational tool. The importance of charge transfer between constituent atom species has been demonstrated as has its inclusion by means of accurate self-consistent solutions. Some examples of the progress achieved in the last year include the following:

A. High Field Superconductivity and Magnetism of Chevrel Phase Compounds

One of the unusual characteristics of the Chevrel phase compounds is that the compounds Mo_6S_8 and Mo_6Se_8 are stable and in the case of Mo_6Se_8 also superconducting with a relatively high transition temperature of 6 K. This transition temperature is comparable to that of PbMo_6Se_8 and LaMo_6S_8 (7K) despite the fact that these compounds have an additional metal atom (Pb, Sn or La).

As proposed, we have studied the electronic structure and properties of Mo_6S_8 and Mo_6Se_8 and, for comparison, LaMo_6S_8 also. As before for the other Chevrels, we have calculated self-consistently the electronic band structure, density of states, electron-phonon coupling parameter and transition temperature for these materials. The results provide us with a first principles understanding of the behavior of these materials and the origin of their superconducting properties. In addition to several publications in international journals, the work has been reported in a book chapter in "Superconductivity in Ternary Compounds". Our work on the Chevrel compounds was presented as an invited paper at the International

Conference on "Superconductivity in d and f Band Metals" in Karlsruhe, Germany, in June, 1982.

B. Superconductivity and Magnetism in C15 Compounds

Our successful studies of the C15 compounds ZrZn_2 and TiBe_2 , discussed in last year's report, led us to undertake an in-depth study of the electronic structure and properties of CeAl_2 and LaAl_2 . These studies were undertaken in order to understand the role played by the 4f electrons - a role obviously essential for explaining the valence fluctuation properties of CeAl_2 . Both paramagnetic and spin-polarized antiferromagnetic state calculations were carried out in order to elucidate the magnetic properties of the system. In the case of CeAl_2 , considerable f electron charge was found on the Ce sites and so justified the use of an extended basis set in the self-consistent procedure to account correctly for the f charge density during the iterative process.

The Ce and La atoms are found to be the dominant factor in determining the electronic structure near the Fermi energy and this is enhanced by the presence of f-bands close to (LaAl_2) or at (CeAl_2) the Fermi energy. In paramagnetic CeAl_2 , the f-bands are about 1 eV wide and, although principally above the Fermi energy, extend down so that the additional electron compared to LaAl_2 is accommodated in the f-density. The ferromagnetic state is found not to be stable but the antiferromagnetic state is found to be stable with the experimentally observed moment. A significant narrowing of the f-bandwidth is observed in the antiferromagnetic state.

C. Soft Phonon Modes and Superconductivity in C15 Compounds

The relationship between electronic, lattice and superconducting properties of materials has attracted considerable theoretical and experimental attention. Recent interest has centered on the C15 super-

conducting compounds HfV_2 and TaV_2 as ideal candidates for these studies. Experimentally, it is known that HfV_2 is a high T_c superconductor ($T_c \sim 9\text{K}$); it has a high critical field, a lattice phase transition at 113 K, a large electronic specific heat coefficient, and a large temperature (T) dependent susceptibility. In contrast, TaV_2 has a low T_c (3.6 K), an average electronic specific heat, an almost T independent susceptibility, and no lattice anomalies. In addition, much is known about certain average properties from a series of heat capacity measurements of $\text{Hf}_{1-x}\text{Ta}_x\text{V}_2$ compounds. These revealed a remarkably strong correlation between the electronic and lattice properties, and showed that the Fermi surface electrons were largely responsible for the soft mode behavior of the phase transition and the high T_c of HfV_2 .

On the theoretical side, our recent detailed self-consistent energy band studies of some Cl5's have elucidated their electronic structure and have yielded some surprising results. For ZrV_2 , the calculations predict a very large electron-phonon coupling $\lambda (=2.4)$ and a very high T_c of 32 K — assuming that the Cl5 structure is retained to low T with unchanged lattice constant and geometric phonon frequencies. These provocative results have raised some questions concerning the role of lattice properties, which are not known but crudely approximated.

Our work has successfully related, for the first time, the microscopic lattice parameters (obtained experimentally) and electronic parameters (obtained theoretically) to the superconductivity of HfV_2 and TaV_2 . First, results of EXAFS measurements taken by Knapp, Georgopoulos and Pan of the mean square relative displacements (MSRD) in HfV_2 and TaV_2 were utilized to calculate some important lattice properties, notably the effective near-neighbor force constants and the effective Debye temperatures of all

pairs of near neighbors. When combined with the results of our self-consistent energy band calculations for HfV_2 and TaV_2 , these constants allowed us to calculate the λ and T_c values of these materials. For TaV_2 , we find $\lambda = 0.4$, consistent with the low T_c , whereas $\lambda = 2.6$ for cubic HfV_2 . Using strong coupling theory, such a large λ would give a T_c of 28 K. This large a value of T_c calculated for HfV_2 is presumably not found experimentally because of the phase transition below which the DOS (and presumably the η values) decrease markedly.

In conclusion, we have shown that the EXAFS technique yields important new information concerning the phonon parameters in a relatively complex system. For the first time all the relevant force constants necessary to calculate T_c have been determined and are found to be much smaller in HfV_2 than in TaV_2 . The increase in the DOS and η values for HfV_2 relative to TaV_2 are responsible for the unusually large decrease in the force constants that drives the lattice phase transition at 113 K.

D. Electronic Structure of TaSe_3 , A Linear Trichalcogenide Compound

Last year we proposed to complete our investigation of the electronic structure and properties of TaSe_3 , a "linear" chain transition metal trichalcogenide compound. The self-consistent electronic energy band structure, density of states, Fermi surface and charge densities have now been accurately determined. These results are now to be compared with experiments being undertaken both at Northwestern University and elsewhere. The calculations represent a forefront state of the art approach to a very complex system of low symmetry (monoclinic structure) which has a total of 16 atoms per unit cell. From the electronic structure results we also hope to understand the origin of the superconducting properties of this system.

E. Assessment of Theoretical Determinations of the Electron-Phonon Coupling Parameter, λ , in Metals and Intermetallic Compounds

The electron-phonon coupling parameter, $\lambda_{\text{el-p}}$, while playing a centrally important role in superconductivity and other phenomena, is still a difficult quantity to determine from first principles theory. The availability of accurate ab initio self-consistent energy band calculations of partial and total density of states (DOS) allow λ to be determined from the electronic specific heat or from simple theoretical treatments such as the rigid ion approximation. We have assessed the accuracy of these determinations (and the band structure results) in a number of transition metals and Al5 and Cl5 intermetallic compounds. We have included comparisons with results obtained using the McMillan equation parameterization of T_c along with experimental results from tunneling measurements, NMR, and comparisons of high and low T specific heat data. As a result of these comparisons, we have been able to show that for many of the high DOS materials serious discrepancies exist between the theoretical and experimental determinations. As a result, the role of spin fluctuations and lattice transformations in connection with these findings has been clarified.

Arthur J. Freeman

Publications supported by AFOSR in 1981-1982

1. "Electronic Structure and Electron Phonon Coupling in Chevrel-Phase Compounds", (with T. Jarlborg) Superconductivity in d- and f-Band Metals 1982, Kernforschungszentrum Karlsruhe 1982.
1. "Assessment of Theoretical Determinations of the Electron-Phonon Coupling Parameter, λ , in Metals and Intermetallic Compounds", (with H.B. Radousky, T. Jarlborg, and G.S. Knapp) Phys. Rev. B 26, 1208 (1982).
2. "Electronic Structure and Itinerant Magnetism in $ZrZn_2$ and $TiBe_2$ ", (with T. Jarlborg and D. D. Koelling) J. Appl. Phys. 52, 1639 (1981).
3. "Electronic Structure and Superconductivity/Magnetism in Ternary Compounds", (with T. Jarlborg) Chapter 6, Vol. 2 of Superconductivity in Ternary Compounds (Springer-Verlag, 1982).
4. "Electronic Structure of Perfect and Defective Solids", TREATISE ON MATERIALS SCIENCE AND TECHNOLOGY (H. Herman, ed.) Vol. 21, Electronic Structure and Properties of Material (F. Fradin, ed.) Academic Press (1981).
5. "Electronic Structure, Spin Polarization and High Critical Fields in Chevrel Compounds" (with T. Jarlborg) J. Magn. and Magn. Matls., 27, 135 (1982).
6. "High-Field Susceptibility and Metamagnetism in Pd" (with T. Jarlborg) Phys. Rev. B 23, 3577 (1981).
7. "Local Density Functional Approach to the Isostructural γ - α Transition in Cerium Using the Self-Consistent Linearized Augmented Plane Wave Method" (with Warren E. Pickett and D.D. Koelling) Phys. Rev. B 23, 1266 (1981).
8. "Self-Consistent Spin Polarized Energy Band Structure and Magnetism in $ZrZn_2$ and $TiBe_2$ ", (with T. Jarlborg and D.D. Koelling). J. Magn. and Magn. Matls. 23, 291 (1981).
9. "Surface Magnetism of Ni Overlayer on a Cu(001) Substrate" (with Ding-sheng Wang and H. Krakauer) Phys. Rev. B 24, 1126 (1981).
10. "Electronic structure and magnetism of CuNi coherent modulated structures", T. Jarlborg and A.J. Freeman, J. Appl. Phys. 52, 1622 (1981).

Arthur J. Freeman

Talks Presented at the Annual Meeting of the American Physical Society,
March 8-12, 1982, Dallas, Texas

1. "Soft Phonon Modes and Superconductivity in C15 Compounds" (G.S. Knapp, H.-K. Pan, A.J. Freeman and T. Jarlborg).
2. "Magnetism versus Superconductivity in HfZn_2 " (Jian-hua Zu, T. Jarlborg, and A.J. Freeman).
3. "Self-consistent Relativistic Electronic Band Structure and Properties of RuO_2 and NbO_2 ", Shi-min Chen, A.J. Freeman, and D. D. Koelling).
4. "Self consistent Relativistic Electronic Band Structures and Properties of A15 Compounds", Shi-min Chen, A.J. Freeman, and D. D. Koelling).
5. "Self-consistent LMO Determination of Electronic Structure of Rutile-dioxides" (Jian-hua Xu, T. Jarlborg and A.J. Freeman).

83
TIC